

■ Psychogenic causes:

Generalized & or Situational:

e.g. psychotic diseases, bad husband and wife relationship, premature ejaculation performance anxiety, homosexuality & other paraphilias ,...etc.

■ *Drugs :*

- Major tranquilizers & most of sedatives and hypnotics,
- Antihypertensive β blockers
e.g, propranolol,
- Diuretics; thiazides,

■ *Chronic diseases :*

- Renal failure(due to hyperprolactinemia).
- Liver disease (↑ estrogen).
- Chronic heart disease.
- Hypertension (arteriosclerosis & atherosclerosis)

■ *iatrogenic causes :*

- Pelvic operations & fracture pelvis: (may affect penile nerves and / or blood supply).
- Total prostatectomy :(nerve affection)

■ *Penile causes :*

- Penile chorde, defective smooth muscles of the cavernous tissue, defective tunical coverings & Pyronie's disease.

2-SPECIFIC TESTS FOR ED .

- **ICI** (intracorporal injection of vasoactive drug)
- **REGISCAN :**
to differentiate between **organic** &
psychogenic ED .
- **CAVOMATE :**
to diagnose **VENO-OCCLUSIVE ED**.
- **DUPLEX :**
to diagnose **ARTERIOGENIC ED** .

Sex Therapy

- It is a couple therapy in the form of Sex education.
- Encourage good relationship between them.
- To get rid off the fear of performance anxiety we advice by Sensate focus program .



Rigiscan

- To measure Nocturnal Penile Tumescence (NPT) & rigidity.
- Normally 3-6 events of erection occur during 8 hours sleep.
- **Absence** or decrease of these events indicates **organic ED**.
- If **normal** it indicates **psychogenic ED**.

Sensate Focus Program

- Instructs the partners to stimulate each other by massage and kissing to gain satisfaction and pleasure and inform each other about the most sensitive and exciting area without genital stimulation.
- Later on, genital stimulation is allowed without intromission. At the end of this program the husband retains his self-confidence

Treatment of E.D.

- **Treatment of the cause if possible**
- **Sex therapy**
- **Medical treatment**
- **Erect Aid Vacuum**
- **Surgical treatment,**
Implantation of penile Prosthesis

Rigiscan

- To measure Nocturnal Penile Tumescence (NPT) & rigidity.
- Normally 3-6 events of erection occur during 8 hours sleep.
- **Absence** or decrease of these events indicates **organic ED**.
- If **normal** it indicates **psychogenic ED**.

■ *Laboratory investigations:*

1-Basic investigations

- *Blood sugar.*
- *Serum prolactin.*
- *Serum testosterone.*
- *Serum lipids*

2-SPECIFIC TESTS FOR ED

Local

1-ICI by self injection of papaverin or prostaglandin before intercourse

NB. If prolonged erection (> 4hours) the patient must contact his doctor to avoid complication

2-Topical application of prostaglandin jelly before intercourse .

3- Transurethral Application

Medical Treatment

Local

1-ICI by self injection of papaverin or prostaglandin before intercourse

NB. If prolonged erection (> 4hours) the patient must contact his doctor to avoid complication

2-Topical application of prostaglandin jell before intercourse .

3- Transurethral Application

Surgical treatment of ED

■ **Penile prostheses implantation.**

■ **Indications:**

- 1-**Severe organic causes**
- 2-**Sever psychogenic causes**
- 3-**when all treatment modalities failed**

SYSTEMIC TREATMENT

- **Phosphodiesterase type 5 Inhibitors**
They are breakthrough treatment for ED
(success rate >70%)
 - Sildenafil (Viagra)
 - Tadalafil (Cialis)
 - Vardenafil (Levitra)

Medical Treatment

LOCAL ttt.

1-ICI

2-Topical application

3- Transurethral Application

SYSTEMIC ttt.

1.Peripheral (PDE5 Inhibitors)

2.Central

3.Both central & peripheral

SYSTEMIC TREATMENT

1. Peripheral (PDE5 Inhibitors)

- Sildenafil (Viagra)
- Tadalafil (Cialis)
- Vardenafil (Levitra)

2. Central

- Dopamine agonist (Apomorphine)

3. Both central & peripheral

- Yohembin
- Melanotan II

Sensate Focus Program

- Instructs the partners to stimulate each other by massage and kissing to gain satisfaction and pleasure and inform each other about the most sensitive and exciting area without genital stimulation.
- Later on, genital stimulation is allowed without intromission. At the end of this program the husband retains his self-confidence

Sensate Focus Program

- Instructs the partners to stimulate each other by massage and kissing to gain satisfaction and pleasure and inform each other about the most sensitive and exciting area without genital stimulation.
- Later on, genital stimulation is allowed without intromission. At the end of this program the husband retains his self-confidence

SYSTEMIC TREATMENT

■ **Phosphodiesterase type 5 Inhibitors**

They are breakthrough treatment for ED

(success rate >70%)

-Sildenafil (Viagra)

-Tadalafil (Cialis)

-Vardenafil (Levitra)

Treatment of the cause

For example :

- Hypogonadism treated with
Androgen replacement.
- Hyperprolactinemia treated with
Bromocriptin.

3-Contraindications

Nitrates

&

Retinitis pigmentosa

(cont.)

- * If refilling or no response to aspiration alone we can aspirate from one corpora & irrigate saline containing V.C. drugs e.g. ephedrine or diluted adrenaline in the other corpora.
(please monitor bl. pressure & heart rate)
- * If detumescence, tight compression & observation for 1 h.
- * If no response shift to the surgical procedures.

Andrology unit

When can you judge on the semen parameters?

- One semen analysis is not enough
- 2 analysis in cases of oligozoospermia
- 3 analysis in cases of azoospermia
- At least 2 weeks between each one
- Recent analysis should be available

HOW TO DEAL WITH **Low-flow** PRIAPISM

- * You must start ttt. as early as possible because tissue ischemia begins after 4-6 hs .
- * Inject mild sedative & pain killer before you start ttt.
- * Encourage ejaculation . If not:
- * Ice packs compresses.
- * **Under complete aseptic condition** ,Aspiration of blood from the corpora cavernosa by 19-gauge scalp needle 20ml by 20ml untile detumesence occurred or for 6 times .

Andrology unit

The color of seminal fluid

- A normal semen has a homogenous, **grayish white** (high protein content)
- **urine in semen**
a **faint yellow** color, & detected by the uriniferous odor
- **Yellow color** in :Jaundice ,carotinemia ,drug
- **Blood in semen** : hematospermia.

Semen

The Semen consists of two components:

1) Seminal plasma

- SV. (65%) -prostate (30%)
- the other accessory glands(5%)

2) Sperms in the seminiferous tubules



Transport of a semen sample to the laboratory:

@ < ½ an hours

@The sample should be protected from extremes of:

- temperature < 20°C & > 40°C
- direct sun rays
- vigorous movement

The viscosity of semen

- The consistency = viscosity & not **coagulation**
- The viscosity measured after liquefaction
- **High viscosity**

associated with:

- **SV hypofunction**
- decreased sperm motility
- infertility (sperm unable to exit from seminal plasma to enter the cervical mucous).

PROGNOSIS OF TTT

The incidence of impotence is directly related to the duration of priapism & the aggressiveness of ttt.

- The ED rate in literature reported to be 50% in low flow priapism ,
- Almost all patients will regain potency if the priapism is aborted within 12-24hs.

Andrology unit

Buried penis in an obese child

- Prepubertal obese Children with a micropenis is caused by the pressure of the prepubic fat on the penis



true penile structure
can be revealed by
pressing the
surrounding fatty
tissue inwards

- The incidence of micropenis in USA IS 15/100.000


(J Urol 2005)

- Micropenis is condition affects 0.6% of men.

Kinesy Institute (2009)

“Trapped penis”

- Penis is trapped after radical circumcision in childhood



Adults Micropenis

- Erect penile length of at least 2.5 standard deviations smaller than the mean penile length

OR

- smaller than about 7 cm ($2\frac{3}{4}$ in) for an adult when compared to an average erection of 12.5 cm (5 in)
- NB.

in a patient with normal internal and external male genitalia.

- USA in the period of 3 years (1997-2000)

The length of the erected penis

- <9cm is 5% of American males
- >20cm is 1% American males
- between 9-20cm is 94% of American males

(J Urol 2005)

Buried penis in obese



Inconspicuous Penis

- This term includes all short penile shaft (micropenis) secondary to abnormalities of the investing structures as in:
 - 1-buried penis in obese
 - 2-trapped penis.
 - 2-webbed penis



(J Urol 1986)

The mean of normal flaccid penis length

- **at birth** is between (2.75 – 4 cm) 4 cm (1.6 in) and the changes very little until puberty, when there is marked growth

BJU International (2007)

- **In adults** is to be 8.9 cm (3.5 inches)

The Journal of Urology (1996)

The Flaccid penis length is a poor predictor of erect length

How to measure penile length



(SPL) Stretched penile length

SPL is measured from the pubis to the tip of the stretched penis



A modified syringe to be used for measuring penile length

- the mean length is approximately 13.12 ± 1.66 cm (5.17 ± 0.65 in).

BJU International (2015)

- the average circumference 12.3 cm (4.8 inches) ,

its average width is 4 cm (1.6 inches)

at N J of Impot Res. (2008)


- the mean length is approximately 13.12 ± 1.66 cm (5.17 ± 0.65 in).

BJU International (2015)

- the average circumference 12.3 cm (4.8 inches),

while its average width is 4 cm (1.5 inches)

Int N J of Impot Res. (2008)



MICROPENIS



**Infantile less
than <2cm**



**Adult smaller
than(<7cm).**

How to measure penile length



(SPL) Stretched penile length

SPL is measured from the pubis to the tip of the stretched penis



A modified syringe to be used for measuring penile length

Normal penile measurements .

- It differs:
 - from country to country &
 - from race to race



Micropenis in a newborn **SPL <2cm**



Normal penile measurements .

- It differs:
 - from country to country &
 - from race to race



The mean of normal erectile measurements


- the mean length is approximately 13.12 ± 1.66 cm (5.17 ± 0.65 in).

BJU International (2015)






- the average circumference 12.3 cm (4.8 inches),

while its average width is 4 cm (1.5 inches)

Int N J of Impot Res. (2008)



Penile development (organogenesis) (8th -12th)

- At the end of 8th week
the bipotential gonadal  **SRY** testis
- Under the effect of maternal
CGH (placenta) on the testis
its Leydig cells  testosterone
its Sertoli cells  MDI
Testo.  **5alpha reductas** DHT  PENILE
differentiated
completely by 12weeks gestation

- Fetal androgen are high between(8-24 weeks)
- Its peak (14-16 weeks)
- Penile growth 20mm from week 16-38
- True micropenis must result from hormonal abnormality that occurs after 12week of gestation

ALSO

- Several homeobox genes defects affect penis and digit size with no detectable hormone abnormalities.
- Intrauterine exposure to certain estrogen drugs like diethylstilbestrol (DES) lead to various genital abnormalities eg. micropenis .
- Exposure to uncontrolled domestic pesticide use increases the likelihood of newborns with a micropenis and/or other genital malformations

TYPES OF MICROPENIS

- CONGENITAL

- 1-Isolated micropenis (**true micropenis**)

Occurs after the 12th week

due to: ↓ gonadotropin → ↓ testosterone

- 2- Micropenis accompanied by
(hypospadias &or undescended testes)
due to: testosterone defect

- Inconspicuous Penis

Penile growth (2nd -3rd trimesters)

- Fetal androgen are high between(8-24 weeks)
- Its peak (14-16 weeks)
- Penile growth 20mm.from week 16-38
- True micropenis must result from hormonal abnormality that occurs after 12week of gestation

Definition

- The term micropenis includes a range of congenital & acquired conditions that result in an abnormally short penis & can be associated with both:
 - functional (related to sex & voiding)
 - psychological problems

Postnatal GROWTH

- Postnatal 1st 6 months fetal pituitary & testes hormones increase → ↑ test. Vol. & penile length
 - The peak of testosterone 1st to 3rd months
 - from (4-6 months) decrease to prepubertal level (Leydig cells) not active
 - So Penile growth during childhood is not androgen dependent
- However the penis doubles in length & diameter between ages of 6 months & puberty

INTRAUTERINE DEVELOPMENT OF THE PENIS

- From 8th to 12th organogenesis
(penil formation)
- During 2nd & 3rd trimesters penile growth

Causes of micropenis

- **1-Insufficient testosterone secretion:**
 - Primary hypogonadism
 - eg. Anorchia
 - klinefelter syndrom
 - Secondary hypogonadism
 - Hypogonadotrophic hypogonadism
 - eg. Kallman syndrome

- **2-Testosterone activation defect**
 - androgen receptors defects
(incomplete form)
 - 5-alpha reductase deficiency
(incomplete form)
- **3-Developmental abnormalities**
eg. penis agenesis
- 4- idiopathic**

Medical ttt.

- Testosterone is effective in treating micropenis due to testosterone deficiency.
- One or two courses of 4 doses testosterone cypionate or enanthate injections
(25 mg) / 3-week
in infancy or childhood resulted in increase in penis sizes to reach the mean for age.
- Topical testosterone application is effective during infancy & childhood (5% cream for one month).

Trapped penis in adult



excessive surrounding



radical circumcision

Trapped penis in adult



excessive surrounding



radical circumcision

Trapped penis(adhesion of penile skin
to the glans)



- Difficulty in :urinating and
:sexual intercourse.
- Some have a low sperm count.
- Psychologically:
 - many have very low self-esteem
 - some even suffer from depression.

HOW TO DEAL WITH MICROPENIS?

- Medical ttt.
- Surgery for micropenis.
- Psychological support

Common clinical presentation

- Most 8- 14 year-old boys referred for micropenis do not have the micropenis . Such conditions are usually explained by one of the following:
 - a penis concealed in suprapubic fat
(Buried & or trapped penis).
 - a large body and frame for which a prepubertal penis simply appears too small.
 - delayed puberty

Micropenis can cause anxiety

- The parents

may worry about future sexual & reproductive functions of their children .

- Young boys

with micropenis might have the fear of discovery by their friends this lead to depression & anxiety

Webbed penis



- The skin connecting the penis to the front side of the scrotum not at the penoscotum

Webbed penis



- The skin connecting the penis to the front side of the scrotum not the penoscotum



Suprapubic fat
pads
surrounding the
penis in the
absence of
additional skin
for the shaft

Suprapubic fat
pads
surrounding the
penis in the
absence of
additional skin
for the shaft

ED = FSAD

■ ED

Persistent inability to attain or maintain

sufficient erection to perform s. intercourse

■ FSAD

Persistent inability to attain or maintain

responses to sexual stimulation

■ Recreational & addictive drugs:

- × **Alcohol:** (liver affection, neuropathy, higher center affection).
- × **Tobacco&smoking:** (asthenosclerosis, vasoconstriction & Decrease of nitric oxide synthetase enzyme.
- × **Opiates& morpheas:** (hyperprolactinemia & C.N.S. affection)
- × **Cannabis:** (C.N.S depression & decrease of testosterone)

■ *Endocrinal causes:*

- Diabetes mellitus (p. neuritis, atherosclerosis)
- Hypogonadism .
- Hyperprolactinemia .
- Hyper & hypothyroidism.

Organic causes (.60%):

■ *Vascular causes e.g.:*

- *Arteriogenic:* (arteriosclerosis, atherosclerosis & stenosis..)

- *Venous leakage ”:*

It may be congenital or acquired abnormalities of the venous drainage

■ ***Neurological causes e.g:***

- Quadriplegia.
- Disseminated sclerosis.
- Temporal lobe epilepsy.
- Spinal cord lesion e.g. Complete sacral injury
(no erection,no ejaculation).

International Society of Impotence Research.

■ Etiology may be:

-Organic > 60%

- Psychogenic < 40%

■ Psychogenic causes:

Generalized & or Situational:

e.g. psychotic diseases, bad husband and wife relationship, premature ejaculation performance anxiety, homosexuality & other paraphilias ,...etc.

CENTRAL NEURAL CONTROL OF EJACULATION

- These central areas mediate the ejaculatory reflex via the interplay between:
- *Primarily:*
 - Serotonergic neurons **Vs** Dopamenergic neurons .
 - (DOPAMIN /SEROTONIN BALANCE THEORY)
 - Waldinger(2002)
- *Secondary:*
 - Cholinergic n.
 - Adrenergic n.
 - GABAergic n.

International Society of Impotence Research.

■ Etiology may be:

-Organic > 60%

- Psychogenic < 40%

■ ***Neurological causes e.g:***

- Quadriplegia.
- Disseminated sclerosis.
- Temporal lobe epilepsy.
- Spinal cord lesion e.g. Complete sacral injury
(no erection,no ejaculation).

■ ***Endocrinal causes:***

■ **Diabetes mellitus (p. neuritis, atherosclerosis)**

■ **Hypogonadism .**

■ **Hyperprolactinemia .**

■ **Hyper & hypothyroidism.**

ED = FSAD

■ ED

Persistent inability to attain or maintain

sufficient erection to perform s. intercourse

■ FSAD

Persistent inability to attain or maintain

responses to sexual stimulation

ED = FSAD

■ ED

Persistent inability to attain or maintain

sufficient erection to perform s. intercourse

■ FSAD

Persistent inability to attain or maintain

responses to sexual stimulation

Sexual Dysfunction (M&F)

■ It is defined as :

impaired or inadequate ability of a man or a women to engage in or enjoy satisfactory sexual intercourse and orgasm

Female Sexual Dysfunctions (FSD)

-- Hypoactive Sexual Desire Disorders

- hypoactive sexual desire**
- sexual aversion**

- Female arousal disorders

- Female orgasmic disorders

- Sexual Pain :

-DYSPAREUNIA
(treatable condition)

-VAGINISMUS

Noradrenaline
Endothelin-1
NPY
PGF_{2α} & TXA₂
Angiotensin II



CONTRACTION

Nitric oxide
Acetylcholine
VIP
PGE₁ & PGE₂
CGRP



RELAXATION


PENILE SMOOTH MUSCLE CELL

Sexual response cycle

(M & F)

- The sexual response cycle (M&F) has the same items and the same definitions for each item:
- 1- Desire
- 2- Arousal
- 3- Orgasm
- 4- Resolution

(American Psychiatric Association ,2000)



Scrotal Swelling Presentations

